

Supplementary Table. Work disability determinants for AAV patients.

Variables	Work disability (n=35)	Preserved work ability (n=53)	Univariate analysis		Multivariate analysis	
			p value	HR (95% CI)	p value	
Demography						
Age, years	46.7 ± 9.5	47.2 ± 12.3	0.38			
Male gender, n	16 (46)	19 (36)				
Disease duration, mo.	116 ± 77	74 ± 57	0.01	1.01 (1.00-1.02)	0.015	
AAV diagnosis						
GPA, n (%)	22 (63)	38 (72)				
MPA, n (%)	1 (3)	5 (9)				
EGPA, n (%)	12 (34)	10 (19)				
Baseline vasculitis involvement						
Constitutional symptoms, n (%)	18 (51)	22/52 (42)	0.51			
Myalgia, n (%)	9 (26)	9/52 (17)	0.42			
Arthralgia, n (%)	13 (37)	16/52 (31)	0.64			
Skin, n (%)	1 (3)	14/52 (27)	0.003	0.78 (0.27-2.26)	0.64	
Eye, n (%)	9 (26)	15/52 (29)	0.81			
ENT, n (%)	29 (83)	39/52 (75)	0.44			
Lungs, n (%)	28 (80)	34/52 (65)	0.16			
Cardiovascular, n (%)	6 (17)	7/52 (13)	0.76			
Gastrointestinal, n (%)	10 (29)	6/52 (12)	0.05	2.49 (0.77-8.02)	0.13	
Kidney, n (%)	11 (31)	11/52 (21)	0.32			
Nervous system, n (%)	14 (40)	17/52 (33)	0.50			
Positive ANCA, n (%)	22 (63)	35/52 (67)	0.82			
Anti-PR3, n (%)	16	23	-			
Anti-MPO, n (%)	4	10	-			
Marital status						
Single	10 (29)	8 (15)		1	-	
Married, divorced or widowed	25 (71)	45 (85)		0.60 (0.20-1.85)	0.38	
Educational level						
No diploma	5 (15)	7 (13)				
Technical degree	10 (30)	13 (25)				
Bachelor degree	5 (15)	7 (13)				
Two-year university degree	5 (15)	4 (8)				
Master degree or more	8 (25)	22 (41)				

*Except where indicated otherwise, values are the number (%) of patients.