

Supplementary Table I. The *MEFV* variants in familial Mediterranean fever (FMF) patients.

<i>MEFV</i> variants	Control FMF patients (n=325), n (%)	Colchicine resistant FMF patients (n=60), n (%)
M694V/ M694V	107 (33)	41 (68.4)
M680I/M680I	18 (5.6)	2 (3.3)
M694V/M680I	33 (10.2)	9 (15)
M694V/V726A	24 (7.4)	2 (3.3)
M694V/ M694I	2 (0.6)	-
M694V/R761H	5 (1.5)	-
M680I/V726A	7 (2.2)	-
M680I/R761H	1 (0.3)	-
E148Q/E148Q	4 (1.3)	1 (1.7)
V726A/F479L	1 (0.3)	-
M694V/E148Q	13 (4)	2 (3.3)
M680I/E148Q	1 (0.3)	-
V726A/P369S	1 (0.3)	-
V726A/E148Q	1 (0.3)	1 (1.7)
E148Q/P369S	2 (0.6)	-
E148Q/R761H	1 (0.3)	-
P369S/P369S	1 (0.3)	-
M694V/-	46 (14.2)	-
M680I/-	10 (3)	-
V726A/-	10 (3)	-
R761H/-	1 (0.3)	-
A744S/-	1 (0.3)	-
E148Q/-	8 (2.5)	-
P369S/-	1 (0.3)	-
F479L/-	1 (0.3)	-
-/-	25 (7.6)	2 (3.3)

MEFV: Mediterranean fever; FMF: familial Mediterranean fever.