

**Supplementary Table I.** Logistic regression analysis on outcome of AOSD patients according to fever pattern.

	Group 2 (n=35)		Group 1 (n=14)		Group 3 (n=21)	
	Odds ratio (95% CI)	<i>p</i> -value	Odds ratio (95% CI)	<i>p</i> -value	Odds ratio (95% CI)	<i>p</i> -value
Systemic score* $\geq 7$	1.00 (reference)	-	4.00 (1.05-15.21)	<b>0.04</b>	1.60 (0.46-5.63)	0.46
Clinical course <sup>†</sup>						
Monocyclic course	1.00 (reference)	-	0.37 (0.10-1.34)	0.13	0.41 (0.14-1.25)	0.12
Polycyclic course	1.00 (reference)	-	2.50 (0.70-8.98)	0.16	2.27 (0.74-7.02)	0.15
Chronic course	1.00 (reference)	-	1.29 (0.21-8.00)	0.78	1.29 (0.26-6.43)	0.76

AOSD, adult-onset Still's disease.

\*Systemic score ranged from 0 to 12, with 1 point awarded for each of the following symptoms: fever, typical rash, sore throat, lymphadenopathy, leukocytosis  $\geq 15,000/\text{mm}^3$ , myalgia, abdominal pain, hepatomegaly, splenomegaly, pneumonia, pleuritis, and pericarditis.

<sup>†</sup>As there was no AOSD-related death in all three group, it was excluded from the analysis.