Supplementary Figure S1. Smoking Habits Questionnaire

SMOKING QUESTIONNAIRE

		1. Visit	Date:/			
Patient's	s initials:					
Date of	Birth:	daymonthyear				
Gender:		•				
Please a	inswer to	the following questions! Circle the answer which a	upplies to you!			
1.	Have you ever smoked a cigarette, cigar or pipe?					
1.	A.	Yes				
	B.	No, I have never tried smoking (end of questions h	ere)			
2.	Have you ever smoked regularly?					
	A.	Yes				
	В.	No, only occasionally				
3.	How old	l were you when you started smoking? years				
4.	What ki	nd of tobacco do you use/did you use?				
	A.	cigarette				
	В.	pipe				
	C.	cigar				
5.	Do you currently smoke?					
	A.	Yes, I smokepieces/day				
	B.	Yes, occasionally				
_	C.	No. (continue with question 8)				
6.		any cigarettes have you smoked recently on average ppieces/day				
7.	Have yo	ou ever tried to stop smoking?				
	A.	Yes, I was not smoking for ye	ars			
	B.	No				
8.	How old	d were you when you stopped smoking?	years			
9.	When did you stop smoking?					
	A.	Within 1 week				
	В.	Within 1–4 weeks				
	C.	Within 1–6 months				
	D.	Within 6–12 months				
	E.	Within 1–5 years				
10	F.	Within more than 5 years	1			
10.		any cigarettes did you smoke on average during yourpacks/day <i>or</i> pieces/day	smoking period?			

Supplementary Table S1. Disease characteristics for all patients from baseline until the end of the study.

Disease activity, mean (SD)	Week 0 9.3 (5.5)	Week 12 1.6 (2.8)	Week 24	Week 52 0.9 (1.9)	Week 104 0.9 (1.9)
Swollen joint count			1.3 (2.5)		
Tender joint count	13.5 (6.4)	2.7 (3.8)	2.2 (3.6)	1.8 (3.6)	1.6 (2.3)
VAS disease assessment by patient (cm)	7.0 (1.8)	3.0 (2.1)	2.5 (2.2)	2.3 (2.1)	2.6 (2.3)
ESR	34.9 (24.5)	21.7 (18.5)	19.1 (17.4)	18.8 (17.4)	15.5 (13.2)
CRP	21.2 (25.2)	10.9 (22.2)	7.6 (14.7)	8.2 (16.5)	5.2 (6.6)

Full analysis set. cm: centimetres; CRP: C-reactive protein; ESR: erythrocyte sedimentation rate; SD: standard deviation; VAS: visual analogue scale.