

Appendix 1

Other members of the Sjögren Big Data Consortium-Sjögren GEAS-SEMI who contributed to this study**Sjögren Big Data Consortium**

- Xiaomei Li, Bei Xu; Anhui Provincial Hospital, China.
- Tamer A. Gheita; Cairo University, Egypt.
- Angelica Gattamelata; Department of Internal Medicine and Medical Specialties, Rheumatology Clinic, Sapienza University of Rome, Italy.
- Alain Sarau, Marie Jezequel; Cavale Blanche-University Hospital, France.

- Béla Nagy; University of Debrecen, Hungary.
- Jorge Sanchez-Guerrero; Instituto Nacional de Ciencias Médicas y Nutrición Salvador Zubirán, Mexico.
- Sarah Downie-Doyle, Chandra Kirana; The Queen Elizabeth Hospital, Adelaide, Australia.
- Gwenny M.P.J. Verstappen; Department of Rheumatology and Clinical Immunology, University of Groningen, University Medical Center Groningen, the Netherlands.
- Arjan Vissink; Department of Oral and Maxillofacial surgery, University of Groningen, University Medical Center Groningen, the Netherlands.

Sjögren GEAS-SEMI

- Miriam Akasbi; Department of Internal Medicine, Hospital Infanta Leonor, Madrid, Spain.
- Richard Alberto Rojas, Anna Maria Febrer Nafria; Hospital Universitari Joan XXIII, Tarragona, Spain.
- Antoni Sisó-Almirall; Primary Care Centre Les Corts, Consorci d'Atenció Primària de Salut Barcelona Esquerra (CAPSBE), Barcelona, Spain.


Supplementary Table S1. Adverse events and disease flares in patients with autoimmune rheumatic disease following SARS-CoV-2 vaccination.

	Machado <i>et al.</i> (1)	Tzioufas <i>et al.</i> (2)	Connolly <i>et al.</i> (3)	Furer <i>et al.</i> (4)	Esquivel- Valerio <i>et al.</i> (5)	Ferri <i>et al.</i> (6)	Sattui <i>et al.</i> (7)	Riberio <i>et al.</i> (8)
Number of vaccinated patients with rheumatic disease	4604	605	1377	686	225	478	2860	910
Types of diseases								
Rheumatoid arthritis, n (%)	1686 (36.6)	168 (27.8)	647(47)	263 (38.3)	132 (58.6)	101 (21.1)	1209 (42.3)	256 (28.1)
Spondyloarthropathies n (%)	1192 (25.9)	126 (20.8)		233 (34)	29 (9.7)	ND	462 (16.2)	195 (21.4)
Systemic lupus erythematosus n (%)	367 (8)	118 (19.5)	273 (20)	101 (14.7)	25 (11.1)	38 (7.9)	391 (13.7)	232 (25.5)
Sjögren's syndrome n (%)	223 (4.8)	58 (9.6)	65 (5)	ND	12 (5.3)	ND	438 (15.3)	42 (4.6)
Vasculitis n (%)	593 (12.9)	67 (11.1)	41 (3)	70 (10.2)	ND	74 (15.5)	167 (5.8)	66 (7.3)
Other diseases n (%)	722 (15.7)	68 (11.2)	351 (23)	19 (2.8)	33 (14.6)	265 (55.4)	929 (32.5)	119 (13.1)
Patients with adverse events n (%)	1894 (41.1)	ND	(D1 87% D2 86%)	ND	173 (76.8)	221 (46.2)	1350 (47.2)	D1 459 (50.5) D2 348 (39)
Total Adverse Event Counts	3334	699	ND		265	287	ND	
Number of local adverse events n (%)	1106 (33.1)	263	ND	D1: 418, D2: 343	158	182	ND	D1 213 (23.4) D2 154 (17.2)
Number of systemic adverse events n (%)	2228 (66.9)	436	ND	D1: 210 D2: 528	107	105	ND	D1 392 (43.3) D2 298 (33.4)
Flares n (%)	204 (3.4)	96 (15.9)	151 (11)	D1: 17 (2.53) D2: 12 (1.79)	ND	ND	382 (13.4)	ND
SARS-CoV-2 infection after vaccination	46 (1)	ND	ND	ND	ND	ND	ND	ND

(D1: after the first dose of the vaccine, D2: after the second dose of the vaccine).

Supplementary Table S2. Demographics and disease-related characteristics of COVID-19-vaccinated patients with pSS.

1. Epidemiological data		3.6. Pulmonary domain	
1.1. Gender		No	1157 (94.1)
F	1170 (94.6)	Low	52 (4.2)
M	67 (5.4)	Moderate	20 (1.6)
1.2. Age at diagnosis (n=1235)	50.5 ± 13.2	3.7. Renal domain	
2. Sicca features at diagnosis		No	1218 (99.1)
2.1. Item I (ocular symptoms)	1141 (92.2)	Low	4 (0.3)
2.2. Item II (oral symptoms)	1134 (91.7)	Moderate	4 (0.3)
Diagnostic tests at diagnosis		High	3 (0.2)
2.3. Item III-1 (Schirmer's test ≤5 mm in 5 minutes in at least one eye)	879/1078 (81.5)	3.8. Muscular domain	
2.4. Item III-2 (ocular staining, ≥4 van Bijsterveld score and/or ≥5 OSS in at least one eye)	328/471 (69.6)	No	1217 (99)
2.5. Item IV (salivary biopsy, focus score ≥1)	675/771 (87.5)	Low	10 (0.8)
2.6. Item V-1 (unstimulated whole salivary flow ≤1.5 mL in 15 min)	582/702 (82.9)	Moderate	2 (0.2)
2.7. Item V-2 (altered parotid sialography)	186/212 (87.7)	3.9. PNS domain	
2.8. Item V-3 (altered salivary scintigraphy)	239/274 (87.2)	No	1175 (95.6)
2.9. Item VI-1 (anti-Ro/SSA)	897/1236 (72.6)	Low	42 (3.4)
2.10. Item VI-2 (anti-La/SSA)	496/1224 (40.5)	Moderate	8 (0.7)
3. Systemic activity at the visit (n=1229)		High	4 (0.3)
ESSDAI domains		3.10. CNS domain	
3.1. Constitutional domain		No	1223 (99.5)
No	1166 (94.9)	Moderate	5 (0.4)
Low	60 (4.9)	High	1 (0.1)
Moderate	3 (0.2)	3.11. Haematological domain	
3.2. Lymphadenopathy domain		No	1094 (89)
No	1122 (91.3)	Low	122 (9.9)
Low	90 (7.3)	Moderate	12 (1)
Moderate	10 (0.8)	High	1 (0.1)
High	7 (0.6)	3.12. Biological domain	
3.3. Glandular domain		No	907/1202 (75.5)
No	1125 (91.5)	Low	207/1202 (17.2)
Low	96 (7.8)	Moderate	88/1202 (7.3)
Moderate	8 (0.7)	4. Type and modification of pre-vaccination SS treatments	
3.4. Articular domain		Treatment modifications	
No	893 (72.7)	No change	1153 (93.8)
Low	257 (20.9)	Delay	3 (0.2)
Moderate	67 (5.5)	Withdraw	68 (5.5)
High	12 (1)	Dose reduction	14 (1.2)
3.5. Cutaneous domain		Treatment types	
No	1178 (95.9)	Biologic treatment	18 (1.5)
Low	21 (1.7)	Glucocorticoids	26 (2.1)
Moderate	28 (2.3)	csDMARDs	28 (2.2)
High	2 (0.2)	No treatment	1147 (93.2)



Supplementary Fig. S1. Distribution of AEs (only local, systemic, and none) stratified according to the ESSDAI activity at the time of the first shot (no activity, low, moderate, and high).

Supplementary Table S3. Patient characteristics associated with sicca exacerbation following vaccination.

Variables	Sicca symptoms worsened		OR [95% CI]	Multivariate OR [95% CI]
	No (n = 1081)	Yes (n = 141)		
Mean age at vaccination	60.6 ± 12.7	59.4 ± 11.8	0.99 [0.98-1.01]	
Age (<60 years)	484/1080 (44.8)	68 (48.2)	1.15 [0.81-1.63]	
Sex (women)	1020 (94.4)	136 (96.5)	1.63 [0.71-4.72]	
Ethnicity (white)	879/1038 (84.7)	121/129 (93.8)	2.74 [1.40-6.19]	1.90 [0.85-4.71]
Country (Europe)	823 (76.1)	124 (87.9)	2.29 [1.39-4.00]	1.64 [0.92-3.14]
Anti-Ro/La (+)	816 (75.5)	91 (64.5)	0.59 [0.41-0.86]	0.58 [0.40-0.86]
Mean ESSDAI at vaccination	3.2 ± 4.6	2.9 ± 4.00	0.98 [0.94-1.02]	
Disease activity state (low)	781/1048 (74.5)	105/140 (75)	1.03 [0.69-1.56]	
Previous SARS CoV2 infection	74/1011 (7.3)	8/140 (5.7)	0.77 [0.33-1.54]	
Vaccine type (RNA)	698 (64.6)	102 (72.3)	1.44 [0.98-2.14]	
Vaccine regimen (incomplete)	86 (8.0)	10 (7.1)	1.13 [0.60-2.37]	

Supplementary Table S4. Characteristics of patients with systemic SS exacerbations in clinical ESSDAI domains after vaccination.

Country	Sex (age)	ESSDAI score at time of vaccination	COVID-19 vaccine	Post-vaccination ESSDAI activity	Exacerbation or new systemic activity	Days after shot	First or second shot	ESSDAI severity	Histopathological study	Treatment	Outcome
Brazil	F (39)	2	nCov-19	Diffuse purpura + Bilateral parotid enlargement	New	7	1	High(cutaneous), Moderate (parotiditis)	ND	Cs	Resolved
Brazil	F (13)	9	nCov-19	Polyarthritits	New	7	1, 2	Moderate	ND	Cs	Resolved
Brazil	F(30)	18	CV	Diffuse purpura	Relapse	10	2	High	ND	Cs	Resolved
Brazil	F(59)	2	BNT	Interstitial lung disease + arthralgia	New (ILD), Relapse (arthritis)	14	1	Low	ND	Cs	Resolved
France	F(57)	2	BNT	Arthritis	New	3	1	Moderate	ND	Cs	Resolved
Italy	F(58)	6	BNT	Diffuse purpura	Relapse	5	1	High	ND	Cs	Resolved
Italy	F(69)	5	BNT	Limited purpura	Relapse	10	2	Moderate	ND	Cs+MTX+ HCQ	Resolved
Slovenia	F(48)	0	BNT	Arthritis	Relapse	5	2	Moderate	ND	Cs+NSAIDs	Resolved
Brazil	F(31)	2	nCov-19	Arthritis, left parotid enlargement	Relapse(arthritis), New (parotiditis)	1	1	Moderate (arthritis), Low (parotiditis)	ND	Cs	Resolved
Brazil	F(42)	0	CV	Limited purpura	New	4	1	Moderate	ND	Cs	Resolved
Netherlands	M(32)	0	BNT	Bilateral parotid enlargement	Relapse	21	2	Low	ND	Symptomatic	Resolved
Netherlands	F(44)	5	BNT	Trigeminal neuropathy	Relapse	21	1	Low	ND	Symptomatic	Resolved
Netherlands	F(42)	ND	BNT	Bilateral parotid enlargement	Relapse	6	2	Low	ND	Symptomatic	Resolved
Netherlands	F(19)	4	BNT	Bilateral parotid enlargement	Relapse	36	2	Low	ND	Symptomatic	Resolved
Netherlands	F(46)	5	BNT	Bilateral parotid enlargement	Relapse	14	1	Low	ND	Symptomatic	Resolved

nCov-19: ChAdOx1 nCov-19; CV: CoronaVac; BNT: BNT162b2; F: female; M: male; ND: not determined; ILD: interstitial lung disease; Cs: corticosteroids; MTX: methotrexate; HCQ: hydroxychloroquine.

References

- MACHADO PM, LAWSON-TOVEYS S, STRANG-FELD A *et al.*: Safety of vaccination against SARS-CoV-2 in people with rheumatic and musculoskeletal diseases: results from the EULAR Coronavirus Vaccine (COVAX) physician-reported registry. *Ann Rheum Dis* 2022; 81: 695-709. <https://doi.org/10.1136/annrheumdis-2021-221490>
- TZIOUFAS AG, BAKASIS A-D, GOULES AV *et al.*: A prospective multicenter study assessing humoral immunogenicity and safety of the mRNA SARS-CoV-2 vaccines in Greek patients with systemic autoimmune and autoinflammatory rheumatic diseases. *J Autoimmun* 2021; 125: 102743. <https://doi.org/10.1016/j.jaut.2021.102743>
- CONNOLLY CM, RUDDY JA, BOYARSKY BJ *et al.*: Disease Flare and Reactogenicity in Patients With Rheumatic and Musculoskeletal Diseases Following Two-Dose SARS-CoV-2 Messenger RNA Vaccination. *Arthritis Rheumatol* 2022; 74: 28-32. <https://doi.org/10.1002/art.41924>
- FURER V, EVIATAR T, ZISMAN D *et al.*: Immunogenicity and safety of the BNT162b2 mRNA COVID-19 vaccine in adult patients with autoimmune inflammatory rheumatic diseases and in the general population: a multicentre study. *Ann Rheum Dis* 2021; 80: 1330-38. <https://doi.org/10.1136/annrheumdis-2021-220647>
- ESQUIVEL-VALERIO JA, SKINNER-TAYLOR CM, MORENO-ARQUIETA IA *et al.*: Adverse events of six COVID-19 vaccines in patients with autoimmune rheumatic diseases: a cross-sectional study. *Rheumatol Int* 2021; 41: 2105-08. <https://doi.org/10.1007/s00296-021-05017-9>
- FERRI C, URSINI F, GRAGNANI L *et al.*: Impaired immunogenicity to COVID-19 vaccines in autoimmune systemic diseases. High prevalence of non-response in different patients' subgroups. *J Autoimmun* 2021; 125: 102744. <https://doi.org/10.1016/j.jaut.2021.102744>
- SATTUI SE, LIEW JW, KENNEDY K *et al.*: Early experience of COVID-19 vaccination in adults with systemic rheumatic diseases: results from the COVID-19 Global Rheumatology Alliance Vaccine Survey. *RMD Open* 2021; 7: e001814. <https://doi.org/10.1136/rmdopen-2021-001814>
- MEDEIROS-RIBEIRO AC, AIKAWA NE, SAAD CG *et al.*: Immunogenicity and safety of the CoronaVac inactivated vaccine in patients with autoimmune rheumatic diseases: a phase 4 trial. *Nature Med* 2021; 27: 1744-51. <https://doi.org/10.1038/s41591-021-01469-5>