

**Supplementary Table S1.** Type of ILD on the CT pattern.

Group	Group A	Group B	Group C	Group D	Group E	All
ILD type on HRCT						
NSIP, n (%)	42 (62.7)	22 (66.7)	12 (46.2)	19 (67.9)	32 (62.7)	127 (62.0)
OP, n (%)	16 (23.9)	8 (24.2)	5 (19.2)	6 (21.4)	6 (11.8)	41 (20.0)
NSIP+OP, n (%)	9 (13.4)	3 (9.1)	9 (34.6)	2 (7.1)	8 (15.7)	31 (15.1)
NSIP+UIP, n (%)	0	0	0	0	2 (3.9)	2 (1.0)
UIP+OP, n (%)	0	0	0	0	2 (3.9)	2 (1.0)
No ILD, n (%)	0	0	0	1 (3.6)	1 (2.0)	2 (1.0)

Group A: glucocorticoid + cyclophosphamide + calcineurin inhibitor; Group B: glucocorticoid + tofacitinib + calcineurin inhibitor; Group C: glucocorticoid + cyclophosphamide; Group D: glucocorticoid + calcineurin inhibitor; Group E: glucocorticoid single or glucocorticoid + other immunosuppressants. ILD: interstitial lung disease; NSIP: nonspecific interstitial pneumonia; OP: organising pneumonia; UIP: usual interstitial pneumonia.