

Supplementary Table S1. Proportion of disease manifestations among all patients, “O” group, and “non-O” group.

	All patients (n=411)	O Group (n=143)	non-O Group (n=268)	<i>p</i> (χ^2)
Mucocutaneous, n (%)	411 (100)	143 (100)	268 (100)	N/A
ROU	408 (99.3)	143 (100)	265 (98.8)	0.555
GU	307 (74.7)	111 (77.6)	196 (73.1)	0.319
PFL	329 (80.0)	115 (80.4)	214 (79.9)	0.891
EN-like lesion	191 (46.5)	63 (44.1)	128 (47.8)	0.473
Pathergy positivity	274 [‡] (77.2)	96 [°] (81.4)	178 [§] (75.1)	0.186
Joint, n (%)	182 (44.3)	65 (45.5)	117 (43.7)	0.727
Peripheral arthritis	169 (41.1)	62 (43.5)	268 (39.9)	0.501
Spondylitis	44 (10.7)	13 (9.1)	31 (11.6)	0.439
Eye, n (%)	139 [°] (33.9)	52 (36.4)	87 [‡] (32.6)	0.441
Uveitis	135 [°] (32.9)	52 (36.4)	83 [‡] (31.1)	0.278
Others	7 [°] (1.7)	2 (1.4)	5 [‡] (1.9)	1.000
Neurologic, n (%)	30 (7.3)	7 (4.9)	23 (8.6)	0.171
Parenchymal	17 (4.1)	4 (2.8)	13 (4.9)	0.319
Vascular	13 (3.2)	3 (2.1)	10 (3.7)	0.556
GIS, n (%)	3 (0.7)	3 (2.1)	0 (0.0)	0.042
Lung, n (%)	8 (1.9)	2 (1.4)	6 (2.2)	0.719
Cardiac, n (%)	3 (0.7)	2 (1.4)	1 (0.4)	0.278
Epididymitis, n (%)	6 (1.5)	0 (0.0)	6 (2.2)	0.096
Others, n (%)	10 (2.4)	4 (2.8)	6 (2.2)	0.744

BD: Behçet's disease; ROU: recurrent oral ulcer; GU: genital ulcer; EN: erythema nodosum; PFL: pseudofollicular lesion; GIS: gastrointestinal system; n: number of patients; N/A: not applicable.

([‡] n=355, [°] n=118, [§] n=237, [°] n=410, [‡] n=267).

Supplementary Table S2. Distribution of overall vascular, venous, and arterial events and their subtypes among all patients, “O” group, and “non-O” group.

	All patients (n=411)	O group (n=143)	non-O group (n=268)	<i>p</i> (χ^2)
Vascular, n (%)	148 (36.0)	39 (27.3)	109 (40.7)	0.007
Venous, n (%)	130 (31.6)	35 (24.5)	95 (35.4)	0.023
ST	28 (6.8)	7 (4.9)	21 (7.8)	0.260
DVT	97 (23.6)	28 (19.6)	69 (25.7)	0.161
IVC	22 (5.4)	2 (1.4)	20 (7.5)	0.009
MV	1 (0.2)	0 (0.0)	1 (0.4)	1.000
PV	4 (1.0)	1 (0.7)	3 (1.1)	1.000
HV	9 (2.2)	1 (0.7)	8 (3.0)	0.171
SVC	8 (1.9)	1 (0.7)	7 (2.6)	0.271
DST	16 (3.9)	3 (2.1)	13 (4.9)	0.169
Others	2 (0.5)	0 (0.0)	2 (0.7)	0.545
Arterial, n (%)	49 (11.9)	11 (7.7)	38 (14.2)	0.053
LE	5 (1.2)	3 (2.1)	2 (0.7)	0.347
UE	1 (0.2)	0 (0.0)	1 (0.4)	1.000
PA	32 (7.8)	7 (4.9)	25 (9.3)	0.110
TA	3 (0.7)	1 (0.7)	2 (0.7)	1.000
AA	12 (2.9)	2 (1.4)	10 (3.7)	0.230
Carotis	2 (0.5)	0 (0.0)	2 (0.7)	0.545
Others	6 (1.5)	3 (2.1)	3 (1.1)	0.423

AA: abdominal aorta; DST: dural sinus thrombosis; DVT: deep vein thrombosis; HV: hepatic vein; IVC: inferior vena cava; LE: lower extremities; MV: mesenteric vein; n: number of patients; PA: pulmonary artery; PV: portal vein; ST: superficial thrombophlebitis; SVC: superior vena cava; TA: thoracic aorta; UE: upper extremities.