Minor salivary gland fibrosis in Sjögren's syndrome is elevated, associated with focus score and not solely a consequence of aging

K.M. Leehan^{1,2}, N.P. Pezant¹, A. Rasmussen¹, K. Grundahl¹, J.S. Moore¹,
L. Radfar³, D.M. Lewis³, D.U. Stone⁴, C.J. Lessard^{1,2}, N.L. Rhodus⁵,
B.M. Segal⁶, R.H. Scofield^{1,7,8}, K.L. Sivils^{1,2}, C. Montgomery¹, A.D. Farris^{1,2}

¹Arthritis and Clinical Immunology Program, Oklahoma Medical Research Foundation (OMRF), Oklahoma City, OK; ²Department of Pathology, University of Oklahoma Health Sciences Center (OUH-SC), Oklahoma City, OK; ³College of Dentistry, OUHSC, Oklahoma City, OK; ⁴Department of Ophthalmology, Johns Hopkins University, Baltimore, MD, and King Khaled Eye Specialist Hospital, Riyadh, KSA; ⁵Division of Oral Medicine and Diagnosis, Department of Diagnostic and Biological Sciences, School of Dentistry, University of Minnesota, MN; ⁶Division of Rheumatic and Autoimmune Diseases, University of Minnesota, MN; 7Department of Medicine, OUHSC, Oklahoma City, OK; ⁸Department of Veteran's Affairs Medical Center, Oklahoma City, OK, USA.

Kerry M. Leehan, MS, PhD Nathan P. Pezant, MS Astrid Rasmussen, MD, PhD Kiely Grundahl, BS Jacen S. Moore, PhD Lida Radfar, DDS David M. Lewis, DDS Donald U. Stone, MD Christopher J. Lessard, PhD Nelson L. Rhodus, DDS, PhD Barbara M. Segal, MD R. Hal Scofield, MD Kathy L. Sivils, PhD Courtney Montgomery, PhD A. Darise Farris, PhD

Please address correspondence to: Dr A. Darise Farris, Arthritis and Clinical Immunology Program, Oklahoma Medical Research Foundation, 825 NE 13th Street, 73104 Oklahoma City, (OK) USA. E-mail: farrisd@omrf.org

Received on February 14, 2017; accepted in revised form on July 10, 2017.

Clin Exp Rheumatol 2018; 36 (Suppl. 112): S80-S88.

© Copyright CLINICAL AND EXPERIMENTAL RHEUMATOLOGY 2018.

Key words: Sjögren's syndrome, minor salivary gland, fibrosis, focus score, aging

For funding information and competing interests, see page S-87.

ABSTRACT

Objective. Evaluate the presence of minor salivary gland (SG) fibrosis in primary Sjögren's syndrome (pSS) as a function of disease pathology or a consequence of ageing.

Methods. Subjects with sicca symptoms attending a Sjögren's research clinic were classified by American European Consensus Group (AECG) criteria as either pSS or non-SS (nSS). Discovery (n=34 pSS, n=28 nSS) and replication (n=35 pSS, n=31 nSS) datasets were evaluated. Minor SG cross-sections from haematoxylin and eosin stained slides were imaged, digitally reconstructed and analysed for percent area fibrosis. Relationships between SG fibrosis, age, and clinical measures were evaluated using Spearman correlations. Association with SS was assessed by: ROC curve, Variable Selection Using Random Forests (VSURF) and uniand bi-variate regression analyses.

Results. SS subjects had significantly more fibrotic tissue in their minor labial salivary glands (median 24.39%, range 5.12-51.67%) than nSS participants (median 16.7%, range 5.97-38.65%, p < 0.0001); age did not differ between groups (average \pm SD pSS 50.2 \pm 13.9 years, nSS 53.8 \pm 12.4 years). In both the discovery and replication data sets, multiple regression models showed that the area of minor salivary gland fibrosis predicted pSS significantly better than age alone. Age-corrected linear regression revealed that the area of minor salivary gland fibrosis positively associated with vanBijsterveld score (p=0.042) and biopsy focus score (p=0.002). ROC curve and VSURF analyses ranked fibrosis as a significantly more important variable for subject discrimination than age.

Conclusion. *SG* fibrosis is an element of pSS pathology that is related to focus score and is not solely attributable to age.

Introduction

Sjögren's syndrome (SS) is a systemic rheumatic autoimmune disorder with cardinal features of chronic, severe dry eyes and mouth and focal lymphocytic infiltrates in salivary and lacrimal gland tissue (1, 2). The aetiology of SS includes genetic risk (3-5), epigenetic (6, 7), environmental (8) and stochastic factors. Causative pathogenic mechanisms remain unclear but involve dysregulation of innate and adaptive immunity (3, 9) and epithelial cell defects (10).

Fibrosis is a common consequence of tissue damage and inflammation (11) and often complicates rheumatic diseases. Several diseases genetically related to SS have well-described fibrotic components, including primary biliary cirrhosis, systemic sclerosis, ulcerative colitis, and systemic lupus erythematosus (12, 13). The presence of autoantibodies (14-18), overactive innate immune pathways such as interferon and NF- κ B (3, 9, 19-21) and tissue inflammation (13, 14, 16, 22-24) are also commonly shared amongst these disorders. Fibrosis in salivary glands (SG) of SS patients has been noted (1, 25-28). However, whether these fibrotic changes merely reflect ageing or are a feature of disease pathology is unclear. Establishment of age-related SG fibrosis in healthy subjects (29) has led to acceptance of SG fibrosis in SS as a consequence of aging and not disease (30-32). Early studies were hampered by the lack of established SS classification criteria. Diagnosis and classification of SS patients relies on a constellation of objective exam results and subjective symptom reporting, yielding a heterogeneous cohort of patients whose individual courses of SS may be dissimilar. Complicating the assessment of fibrosis is the late age of onset of the disorder, usually in the fourth decade or later. A recent study involving 264 subjects with pSS found that fibrosis grade associated positively with dental damage and was inversely correlated with salivary flow (27). Another study evaluating SG biopsy slides of 63 cases of pSS and 11 healthy controls reported an increased grade of fibrosis in pSS and an association between fibrosis and SS after adjusting for age (28). While these studies have subjectively (using graded scales (25, 29, 33)) assessed fibrosis, no study has quantified fibrosis in the minor SGs in order to dissect the relationships between fibrosis, age and SS. Determining whether fibrosis is a form of pathology in SS contributes to our overall understanding and may open new therapeutic avenues for SS patients.

This study was undertaken to determine if SS subjects have more, less, or similar fibrotic replacement as compared to subjects who have symptoms of dryness but do not meet established disease criteria (non-SS) and to determine whether the presence of fibrosis is solely attributable to age. We report that SG fibrosis is a pathologic feature of SS related to focal SG inflammation and not solely a consequence of age. An age-related increase in minor SG fibrosis is confirmed and this study further establishes that minor SG fibrosis is an element of lymphocytic focus-associated SG pathology and is not solely an attribute of older subjects.

Participants and methods

Participants

Biological samples, clinical and laboratory test values were obtained from the Sjögren's Research Clinic (SRC) (at Oklahoma Medical Research Foundation and University of Minnesota) as previously described (34). Clinical measures including focus score, van Bijsterveld score, Schirmer's test, and collection of whole unstimulated salivary flow, were conducted as specified by the 2002 revised American European Consensus Groups (22). Ro antibody titres were determined as described (35, 36). Briefly, IgG antibody titres were determined by ELISA using bovinederived Ro60 (Immunovision).

The participants were self- or physician-referred, underwent pre-clinical Table I. Subject demographic characteristics.

| | Discovery Set | | | Replication Set | | |
|-------------------------------------|---------------|---------|----------|-----------------|---------|----------|
| | pSS | DNMC | p-value* | pSS | DNMC | p-value* |
| Total participants (n) | 34 | 28 | | 35 | 31 | |
| Age mean (SD) | 52.38 | 51.82 | 0.8713 | 56.20 | 50.58 | 0.833 |
| | (12.34) | (14.38) | | (11.49) | (14.08) | |
| Fibrosis mean (SD) | 23.05 | 15.24 | 0.0002 | 26.85 | 18.52 | 0.0002 |
| | (8.63) | (6.02) | | (10.32) | (7.28) | |
| Female (%) | 82.35 | 85.71 | 1 | 88.57 | 90.32 | 1 |
| Caucasian (%) | 100 | 89.29 | 0.0866 | 85.71 | 64.52 | 0.0829 |
| Anti-Ro/SS-A positive (%) | 64.7 | 0 | <0.0001 | 65.71 | 3.22 | <0.0001 |
| Anti-La/SS-B positive (%) | 47.06 | 3.57 | 0.0001 | 34.26 | 6.45 | 0.0068 |
| WUSF positive (%) | 64.71 | 46.43 | 0.2 | 62.85 | 29.03 | 0.0074 |
| Schirmer's positive (%) | 55.88 | 35.71 | 0.1323 | 45.71 | 29.03 | 0.2073 |
| VanBijsterveld positive (%) | 61.76 | 35.71 | 0.0403 | 57.14 | 22.58 | 0.0059 |
| FS≥1 (%) | 73.53 | 17.24 | <0.0001 | 54.29 | 12.9 | 0.0006 |
| Anti-cholinergic drugs positive (%) | 52.94 | 67.86 | 0.3012 | 62.86 | 48.39 | 0.3212 |

**p*-values were calculated by Fisher's exact test, except for age (unpaired 2-tailed *t*-test), and fibrosis mean (unpaired 2-tailed Mann-Whitney test).

screening using questions pertaining to oral and ocular disease symptoms (35) and had at least one qualifying ocular and one qualifying oral dryness complaint. All participants gave fully informed consent in compliance with the Declaration of Helsinki, and the study was approved by both respective Institutional Review Boards. Participants were classified using the 2002 revised AECG criteria (22).

All participants with features of overlapping diseases (including rheumatoid arthritis, systemic lupus erythematosus, and systemic sclerosis) or with exclusion criteria for AECG classification (sarcoidosis, prior head and neck radiation, hepatitis C infection, acquired immunodeficiency syndrome, pre-existing lymphoma, and graft-versus-host-disease) were excluded from the study. Participants who failed to meet AECG criteria for pSS, but had dry eye and/or dry mouth complaints were designated as "non-SS" (nSS). Participants in pSS and nSS categories were randomised into separate datasets, a discovery set (n=34 pSS, n=28 nSS) and a replication set (n=35 pSS, n=31 nSS). Randomisation was performed using discrete uniform distribution sampling via the "sample" function in R. Imaging and fibrosis scoring of participants' biopsy tissue was performed with classification status blinded. Demographic and clinical data of participants are shown in Table I. Dental data, including number of tooth restorations, was available for a subset of the subjects (n=44 pSS, n=36 nSS). Disease duration data was abstracted from patient questionnaires, and represent the most conservative estimation based on the date of diagnosis (age of study entry) and the calculated age of symptom onset (based on questions regarding subjective dryness).

Salivary gland biopsy and imaging

Four to six minor labial SGs per participant were formalin-fixed, paraffin-embedded, sectioned (4 µm), and stained with haematoxylin and eosin in either the University of Minnesota or University of Oklahoma Health Sciences Center oral pathology laboratories. Focus scores were determined by a boardcertified oral pathologist. The slide used for focus scoring was imaged using a Zeiss 710 confocal microscope. Each glandular cross-section was imaged at 200x magnification in overlapping sections. These sections were digitally assembled using the Zeiss ZenBlue software package to yield reconstructions of entire glandular cross-sections (Fig. 1).

Quantitative fibrosis assessment

Fibrotic changes were quantified by an observer blinded to disease classification status as described (36). A standard grid of 2500 μ m² was applied to SG cross-section images using ImageJ (National Institutes of Health, Bethesda, MD, USA). Each individual square



Fig. 1. Scoring of minor labial salivary gland fibrosis area. **A:** Digital reconstruction of a minor labial salivary gland paraffin-embedded tissue section. Boxed region is magnified in B. **B:** Representative scoring of labial SG tissue. "0" indicates squares where tissue is present, but is non-fibrotic. "1" indicates both tissue presence and fibrosis. Magnification = 200 x scale = $1.54 \text{ pixels/}\mu\text{m}$, grid square = $2500 \mu\text{m}^2$.

of each section was scored using the following rubric: edges were accounted for by omitting any square where less than 50% of the area contained tissue (Fig. 1). Areas of infiltration were included in the total area calculation but were assigned a value of '0'. Tissue positive squares containing $\geq 50\%$ fibrotic tissue were assigned a value of 1. Tissue positive squares containing <50% fibrotic tissue were assigned a value of 0. The number of fibrosis positive squares in each cross-section was multiplied by 2500 μ m² (area of each grid square). This value was divided by the total section area to generate the percent area fibrosis for each glandular cross-section for each participant:

 $\left\{\frac{(n=2500um^2)}{section\ area}\right\} \ge 100 = \% \ area \ fibrosis$ (where $n = number\ of\ fibrosis\ positive\ grid\ squares$)

The individual cross-section percent areas were then averaged to yield a perparticipant mean percent-area fibrosis of minor labial SG tissue:

$$\left\{ \frac{(\sec 1\% \text{ area}) + (\sec 2\% \text{ area})...(\sec x\% \text{ area})}{total section number} \right\}$$

= average % area fibrosis

Fibrosis severity scores

A board-certified oral pathologist was provided with SG slides from a selected subset (pSS=20, nSS n=15) of subjects to independently assess degree of fibrosis. The observer was blinded to disease classification of the subject samples. Distribution analysis of fibrotic area was used to select SG slides for independent evaluation; slides were selected from each 'bin' equal to: $\leq 5\%$, 6-20%, 21-30%, and $\geq 30\%$ to cover the full data range. Slides were scored as follows: 0=normal tissue up to very minor periductal fibrosis; 1=significant periductal fibrosis only, 2=acinar replacement by fibrotic tissue with periductal fibrosis, 3=widespread fibrosis including acinar replacement, lobular dysmorphia and extensive gland disruption.

Statistical analyses

All were executed in R (37) or Prism 6.0 (GraphPad Software, La Jolla California USA, www.graphpad.com). Normality tests: two-tailed Shapiro-Wilks tests and where necessary, nonparametric tests. Both bivariate and univariate logistic regression generalised linear models (GLMs) were performed to assess association of fibrosis with disease. Simple linear regressions as well as linear regressions with variable correction (to assess association of fibrosis with clinical measures) utilised Box-Cox transformed data (powerTransform and bcPower functions in the 'car' R package (38)). ROC curves were generated using the R package 'pROC' (39); DeLong's test was used to measure likeness of ROC curves for fibrosis, focus score and age. Maximal Youden's index, as determined by the 'OptimalCutpoints' package (40) was used to determine the optimal threshold for fibrosis in predicting diagnosis. For random forest analysis, the default 'VSURF' (41) package was used to test the importance of average percent area fibrosis and age as well as categorical variables of sex, race, and AECG-determined diagnostic cutoffs for positivity of the following parameters: AECG questions on oral symptoms (yes/no), AECG questions on ocular symptoms (yes/no), vanBijsterveld score \geq 4, Schirmer's score \leq 5 mm/minute, whole unstimulated saliva flow \leq 1.5 mL in 15 minutes, focus score \geq 1, presence of IgG anti-Ro/SSA or IgG anti-La/SSB.

Results

Fibrosis is elevated in the minor labial salivary glands of subjects with primary Sjögren's syndrome To assess the presence and extent of fibrosis in minor salivary glands, a precise method of assessing fibrosis in haematoxylin and eosin stained SG biopsy tissue sections was implemented. Percent area fibrosis values are reported as the average of multiple (4 to 6) glandular cross-sections per subject (n=128 subjects). Average percent area fibrosis in the discovery and replication sets yielded similar results, with SG fibrosis being significantly greater in subjects with pSS compared to those in the nSS group (Fig. 2A & C). In the combined dataset, pSS participants had a greater median percent area SG fibrosis (24.39%, range 5.12-51.67%) than nSS participants Fig. 2. Salivary gland fibrosis but not age is increased in pSS compared to nSS subject groups. A: Average percent area fibrosis is significantly greater in pSS subjects (23.6%±1.1; mean±SEM) as compared to nSS subjects (16.6%±0.90). Mann-Whitney *t*-test.

B: pSS subjects (53.8 ± 1.37) years; mean±SD) are not significantly different in age as compared to nSS subjects (50.2 ± 1.76) years). Student's two-tailed unpaired *t*-test.

C-D: Distribution of percent area fibrosis and subject age are shown.



(16.7%, range 5.97–38.65%, *p*<0.0001). Importantly, there was no significant difference in participant age between pSS and nSS groups (Fig. 2B & D). As expected, the pSS group exhibited higher incidence of positivity for measures utilised to classify individuals as having SS, including positivity for IgG anti-Ro/ SS-A and anti-La/SS-B antibodies, van-Bijsterveld score (ocular damage) and biopsy focus score (focal lymphocytic infiltration) (Table I). In contrast, the percentage of pSS versus nSS subjects with a positive Schirmer's test, indicating reduced tear flow, was not different in either the discovery or replication sets, while the fraction of subjects with a positive whole unstimulated salivary flow (WUSF) test, indicating reduced salivary flow, was only significantly different in the replication set.

To compare the percent area fibrosis measurement with pathologist-determined severity of SG fibrosis, an oral pathologist evaluated tissue slides from a sample of the subjects. This analysis utilised slides from a subset (n=35 subjects) of subjects evaluated for average percent area SG fibrosis measurements covering the range of quantitative fibrosis observed. The pathology scores, which took into account proximity of fibrosis to ducts or acini and related acinar cell destruction, correlated significantly (r=0.6, p=0.0002) with the quantitative data (Fig. 3), indicating that the quantitative percent area fibrosis measurement captured the severity of fibrotic changes. Additionally, a Mann-Whitney U-test comparing the pathologist-assigned scores showed significantly greater fibrosis severity in pSS as compared to nSS groups (p=0.04). Thus, the subjective fibrosis severity score correlates positively with our quantitative measure of fibrosis and recapitulates the elevated fibrosis phenotype observed in pSS subjects.

Fibrosis discriminates pSS from non-SS sicca more effectively than age in regression models

In order to exclude age as a confounding covariate in the assessment of fibrosis in SS, a two variable logistic regression model was employed. SS disease status was the dependent variable, with age and percent area fibrosis as the predictive variables (Table II). In both the discovery and replication analyses, fibrosis contributed to prediction of pSS classification (discovery set p=0.0009, OR=1.16, accuracy=68%; replication set p=0.0060, OR=1.12, accuracy=73%) while age did not (discovery set p=0.5720, OR=0.99, replication set p=0.8346, OR=1.00).

To compare the efficacy of age or fibrosis alone to distinguish pSS versus nSS, single variable models using only fibrosis or only age were constructed and compared to a multivariate model. ANOVA analysis revealed that only fibrosis significantly discriminated between pSS and nSS (discovery set p=0.0010, OR=1.15, accuracy = 69.4%, replication set p=0.0021, OR=1.12, accuracy = 73.0%). Finally, we compared the individual models to the full model. and found that the fibrosis-only model was not significantly different from the multivariate model in distinguishing pSS versus nSS status (discovery set p=0.5695, replication set p=0.8348), indicating that the addition of age does not significantly improve the fibrosis model. Within the multivariate models, **Fig. 3.** Significant agreement between fibrosis severity scores and quantified percent fibrosis by area (pSS=20, nSS n=15), 2-tailed Spearman correlation.



 Table II. Fibrosis contributes significantly more information to discrimination models than age.

| Set* | #Variables | | Point Estimate | <i>p</i> -value (z) | OR | Accuracy | Sensitivity | Specificity | p-value+ |
|------|------------|----------|-------------------|---------------------|------|----------|-------------|-------------|----------|
| DS | 2 | Fibrosis | 0.145 | 0.001 | 1.16 | 0.677 | 0.643 | 0.706 | N/A |
| | | Age | -0.013 | 0.572 | 0.99 | | | | |
| | 1 | Fibrosis | 0.140 | 0.001 | 1.15 | 0.694 | 0.679 | 0.706 | 0.570 |
| | 1 | Age | 0.003 | 0.867 | 1.00 | 0.548 | 0.000 | 1.000 | <0.0001 |
| RS | 2 | Fibrosis | 0.113 | 0.006 | 1.12 | 0.727 | 0.677 | 0.771 | N/A |
| | | Age | 0.005 | 0.835 | 1.01 | | | | |
| | 1 | Fibrosis | 0.116 | 0.002 | 1.12 | 0.727 | 0.677 | 0.771 | 0.835 |
| | 1 | Age | 0.035 | 0.083 | 1.04 | 0.682 | 0.581 | 0.771 | <0.0001 |

*DS: Discovery Set; RS: Replication Set. *ANOVA, as compared to multivariate model.

Table III. Focus score and fibrosis combined enhance precision.

| Set* | | [#] Variables | Point estimate | <i>p</i> -value | OR (z) | Accuracy | Sensitivity | Specificity | <i>p</i> -value ⁺ |
|------|---|-------------------------|-------------------|-----------------------|--------------|----------|-------------|-------------|------------------------------|
| DS | 2 | Focus score Fibrosis | 1.287 | 0.011 0.055 | 3.62 1.10 | 0.790 | 0.857 | 0.735 | N/A |
| | 1 | Focus score | 1.384 | 0.003 | 3.99 | 0.774 | 0.821 | 0.735 | 0.039 |
| | 1 | Fibrosis | 0.140 | 0.001 | 1.15 | 0.694 | 0.679 | 0.706 | <0.0001 |
| RS | 2 | Focus score Fibrosis | 0.674 0.112 | 0.021 0.007 | 1.96 1.12 | 0.758 | 0.807 | 0.714 | N/A |
| | 1 | Focus score | 0.869 | 0.010 | 2.38 | 0.682 | 0.839 | 0.543 | 0.002 |
| | 1 | Fibrosis | 0.116 | 0.002 | 1.12 | 0.727 | 0.677 | 0.771 | 0.0002 |

Table IV. Association of fibrosis with SS clinical features.

| | Linear r | egression | Age-corrected linear regression | | |
|----------------------|-------------------|-----------------|---------------------------------|-----------------|--|
| Clinical variable | Beta | <i>p</i> -value | Beta | <i>p</i> -value | |
| Age | 0.034 | <0.0001 | NA | NA | |
| vanBijsterveld score | 0.174 | 0.028 | 0.152 | 0.042 | |
| Schirmer's score | -0.104 | 0.093 | -0.054 | 0.366 | |
| WUSF | -0.426 | 0.025 | -0.145 | 0.466 | |
| Biopsy focus score | 0.836 | <0.0001 | 0.713 | 0.002 | |
| Dental restorations | 0.083 | 0.012 | 0.042 | 0.245 | |
| WUSE: Whole unstimul | ated salivary flo | W | | | |

age was not a significant contributor to model outcome (discovery set p=0.867, accuracy 54.8%, OR=1.1, replication set p=0.083, accuracy=68%, OR=1.0,). Additionally, the age-only model was inferior to both the fibrosis-alone and multivariate models, indicating that the addition of the fibrosis measure significantly improves the age-only model (ANOVA vs. full model discovery set *p*<0.0001, replication set *p*<0.0001). Taken together, these results indicate that the effect of fibrosis is separate from that of age. This supports the hypothesis that the elevated fibrosis observed in SS patients is not solely attributable to advanced patient age.

Regression modelling reveals that the addition of fibrosis to focus score more precisely discriminates between pSS and non-SS sicca subjects

Biopsy focus score is a benchmark feature used to classify SS. To determine whether fibrosis can enhance the power of focus score in discriminating pSS from nSS subjects, regression modelling was employed. Separation of pSS from nSS was the dependent variable, while biopsy focus score and percent area fibrosis were the predictive variables (Table III). In the discovery set, focus score significantly contributed to SS discrimination (p=0.011) while fibrosis exhibited a trend toward significant contribution to categorisation (p=0.055). However, in the replication set both focus score (p=0.021) and fibrosis (p=0.0067) had the capacity to distinguish between pSS and nSS. This multivariate model was compared to univariate logistic regression models containing only focus score or only fibrosis. In both sets, the multivariate model was significantly better at discriminating the disease groups than either univariate model, indicating that the inclusion of average percent area fibrosis enhances the power of focus score alone (discovery set p < 0.0391, replication set p < 0.002).

Minor salivary gland fibrosis associates with focus score,

ocular damage and age

The relationship between percent area fibrosis, age and other clinical features

was evaluated. Using simple linear regression, age (p<0.0001), biopsy focus score (p<0.0001), vanBijsterveld score (*p*=0.028), WUSF volume (*p*=0.025), and number of tooth restorations (p=0.012), were found to significantly associate with percent area fibrosis (Table IV). Anti-Ro autoantibody titre was evaluated within the pSS population for correlation with fibrosis, and no significant relationship was discovered (p=0.30, Spearman 2-tailed t-test). We also compared the extent of fibrosis in Ro-positive and Ro-negative subjects, but found no significant difference between the groups (Ro+ primary=44, Ro⁻ primary=24, p=0.23, Kolmogorov-Smirnov 2-tailed test.) Interestingly, we found no correlation between patientreported duration of disease and extent of fibrosis. A subset of primary patients (n=52) had disease duration data available, but no association with the degree of fibrosis was apparent (p=0.67, r=0.06). As age is a potentially confounding factor in these analyses, we age-corrected the linear regressions comparing fibrosis and clinical SS signs. Only vanBijsterveld score (p=0.042) and biopsy focus score (p=0.002) associated with degree of SG fibrosis (Table IV). Thus, while age and fibrosis are correlated, fibrosis is correlated with vanBijsterveld score and focus score beyond the contribution of age.

Fibrosis discriminates pSS from non-SS sicca more effectively than age by receiver operating characteristic (ROC) and random forest analyses

As univariate regression analyses showed that focus score and age are most closely associated with fibrosis, we directly compared the capacity of these factors to distinguish between the pSS and nSS groups by ROC analysis. Areas under the ROC curves (AUC) for age, average percent area fibrosis, and biopsy focus score were 57.58, 75.84, and 77.73%, respectively (combined dataset). DeLong's test for two correlated ROC curves showed a significant difference between fibrosis and age (p=0.0012), but not between fibrosis and focus score (p=0.7292) (Fig. 4). We chose a threshold of 20.69% fibrosis to classify subjects because it optimises



Fig. 4. Receiver operating characteristic (ROC) curves for age (black), average percent area fibrosis (red), and biopsy focus score (blue) are shown compared to the line of no-discrimination (dashed blue line). The optimal cut-point distinguishing pSS from DNMC is indicated by the intersection of the dashed red lines.

both specificity (0.780 (0.653–0.877 95% CI)) and sensitivity (0.710 (0.588–0.813 95% CI)).

To compare the capacity of SG fibrosis to distinguish pSS from nSS subjects with other tests used in pSS classification, we used a non-parametric method of variable ranking and selection (by way of random forests). Via an iterative process, variables that do not contribute to the output are eliminated. Continuous variables were limited to average percent area fibrosis and subject age. Categorical variables included sex, race, AECGquestions regarding dry eye and mouth (at the time of clinic visit), as well as the results of objective pSS classification tests including highest vanBijsterveld score, lowest Schirmer's value, WUSF test, anti-Ro/SS-A and anti-La status/ SS-B, and lip biopsy focus score. Only five of these twelve variables passed the importance threshold; they are, in decreasing order of importance, anti-Ro/ SS-A positivity (importance 0.1639), biopsy focus score ≥1.0 (0.0838), anti-La/ SS-B positivity (0.0587), average percent area fibrosis (0.0284), and WUSF \leq 1.5 ml/min (0.0144). Thus, degree of SG fibrosis selectively associates with the SS disease state, whereas subject age exerts no influence on SS disease state by random forest analysis.

Discussion

This study is the first to quantitatively evaluate minor salivary gland fibrosis in subjects with pSS compared to those with sicca symptoms alone. In our analyses, fibrosis distinguishes pSS from those with sicca symptoms who do not meet criteria for SS and performs comparably to biopsy focus score in this regard. As our analyses did not include individuals meeting criteria for other rheumatic diseases, we have not evaluated the extent of salivary gland fibrosis as a tool for disease classification or diagnosis; rather, we offer compelling evidence that fibrosis is part of the SS disease process and not only a consequence of aging.

Our method considered and assessed all fibrotic tissue without prior knowledge of subject classification. High positive correlation of our quantitative measurements with the pathologist-assigned fibrosis severity scores demonstrates that the quantitative method captures changes considered to be of pathologic importance. We report here that subjects classified as pSS have higher average percent area fibrosis than those who do not fulfill pSS classification criteria. Notably, there was no significant difference in age between pSS and non-SS sicca groups in either dataset. Across the entirety of data available from our research centre, however, we observe a significant difference in age between pSS (pSS n=635, median=56 and nSS groups (median=52, n=765, p < 0.0001) (unpublished data). In light of this difference, seen in the larger sample set, we treated age as an independent variable in our analyses, to avoid its confounding effects.

We tested whether fibrosis would associate with other clinical features of SS and detected positive relationships between biopsy focus score and SG fibrosis, and between ocular surface damage (vanBijsterveld score) and fibrosis. The inverse relationships between fibrosis and tooth restorations and fibrosis and WUSF were explained by age, and showed no significant association after age correction. Using multiple approaches, we dissected the individual contributions of age, fibrosis and biopsy focus score in separating subjects with pSS from those in the nSS group. We compared the ability of quantified fibrosis to discriminate between pSS and non-SS nSS groups, as compared to that of age and focus score. We took a threefold statistical approach to better elucidate potential relationships between these three variables: multivariate regression modelling, ROC curve comparison, and random forest variable ranking.

The results showed that, although age and fibrosis correlate, age alone could not explain the extent of fibrosis in pSS subjects as compared to their similarlyaged nSS counterparts. By comparing uni-to bi-variate regression models, we demonstrated that the addition of fibrosis significantly improves the age-alone model and increases the sensitivity and accuracy of the focus score model. These data strongly suggest an intimate relationship between lymphocytic infiltration and fibrotic tissue replacement. The present study undoubtedly underestimates this relationship since SG foci by definition (42) and according to current classification criteria (22) must not be adjacent to fibrotic tissue.

ROC analysis demonstrated that fibrosis out-performed age in predicting pSS versus nSS. In fact, fibrosis was comparable to the predictive power of biopsy focus score in this analysis. A relationship between degree of SG fibrosis and SG lymphocytic infiltration is further supported by our recent observation that the degree of SG CD4+ T cell clonal expansion positively correlates with percent area SG fibrosis in pSS (36). The random forest approach identified the five most important variables as anti-Ro/SS-A positivity > biopsy focus score > anti-La/SS-B positivity > extent of fibrosis > loss of saliva secretion (WUSF). The results of these analyses agree that fibrosis is a variable of importance in stratifying SS versus nSS subjects. While the exact cause of SG fibrosis and its role in SG dysfunction and SS disease remain unknown, the data presented here establish fibrosis as a pathologic feature of SS. The multimodel approach confirms that fibrosis makes a significant contribution to distinguishing non-SS sicca from SS, and that it does so above and independent of the contribution of age. These results are in agreement with those of Llamas-Gutierrez et al. (28), who observed an age-independent association between grade of fibrosis and pSS but included only 11 non-SS controls. Our results present strong, replicated evidence that quantified fibrosis is a feature of Sjögren's syndrome pathology and is not solely a feature of age.

Tissue fibrosis is a common consequence of chronic inflammation, suggesting that the theory of SG fibrosis in SS is plausible, if not probable. CD4⁺ T cells, macrophages and epithelial cells all play roles in both normal homeostasis and pathological accumulation of collagen (43, 44) and are commonly found in glandular lesions in SS (25, 30, 31, 45-47). Increased fibrotic change is correlated with the presence and degree of CD4⁺ T cell clonal expansions in the minor salivary glands (36). Moreover, diseases sharing genetic overlap with primary SS include inflammation-associated tissue fibrosis as a pathological feature (12, 48). For example, in systemic sclerosis, genetic variants of *STAT4* and *IRF5*, (which associate with pSS (48)), demonstrated additive effects contributing to interstitial lung disease (49).

One of the confounding factors in SS disease research is the near-total lack of longitudinal data and the difficulty in precisely capturing theoretical disease duration from self-reported patient data. Disease duration, as it relates to SS, is a difficult parameter to capture, as 1) the onset of irritating dry eye and mouth is difficult to pinpoint in hindsight, and 2) when asked in different ways, patient's responses can be inconsistent. In our study, limited data on patient-reported disease duration was available, and no correlation between disease duration and the extent of salivary gland fibrosis was detected. Although fibrosis is widely considered to be a progressive process, it is possible that salivary gland fibrosis in SS is not progressive. Kapsogeorgou et al. detected no fibrotic progression in labial salivary gland biopsies longitudinally collected a median of 4.5 years apart (50). We also note that patient-reported disease duration is an imprecise measure.

Determining the disease chronology and sequence of events leading to glandular hypofunction in SS can only be accomplished by well-designed and comprehensive longitudinal studies. Recognising lymphocytic focus-associated SG fibrosis as a fundamental pathology in SS, however, furthers our understanding of the complexity of this disease and paves the way for future investigations evaluating the utility of this feature for sub-setting SS patients.

Acknowledgments

The authors gratefully acknowledge the contributions of Michael Brown, Ben Fowler, Justin Willige and the OMRF Imaging Core facility for digital recon-

struction of tissue cross-sections, Dr Rajaram Gopalakrishnan, Dr Andrew J.W. Huang, Dr Pamela J. Hughes, and Dr Michael D. Rohrer for their contributions to the University of Minnesota SRC.

Funding

This work was supported by NIH grants AR060804, AR050782, DE015223 and DE018209, COBRE GM110766, Phileona Foundation, and by an American Association of Dental Research Sjögren's Syndrome Foundation Student Fellowship to K.M. Leehan.

Competing interests

R. Hal Scofield has received grant support for research related to Sjögren's syndrome from the United States government; the other co-authors have declared no competing interests.

References

- JONSSON R, VOGELSANG P, VOLCHENKOV R, ESPINOSA A, WAHREN-HERLENIUS M, APPEL S: The complexity of Sjögren's syndrome: Novel aspects on pathogenesis. *Immunol Lett* 2011; 1: 1-9.
- 2. MAVRAGANI CP, MOUTSOPOULOS HM: Sjögren's syndrome. *Annu Rev Pathol* 2014; 273-85.
- LESSARD CJ, LI H, ADRIANTO I et al.: Variants at multiple loci implicated in both innate and adaptive immune responses are associated with Sjögren's syndrome. Nat Genet 2013; 11: 1284-92.
- NOCTURNE G, MARIETTE X: Advances in understanding the pathogenesis of primary Sjögren's syndrome. *Nat Rev Rheumatol* 2013; 9: 544-56.
- LIY, ZHANG K, CHEN H et al.: A genome-wide association study in han chinese identifies a susceptibility locus for primary Sjögren's syndrome at 7q11.23. Nat Genet 2013; 11: 1361-5.
- ALTOROK N, COIT P, HUGHES T et al.: Genome-wide DNA methylation patterns in naïve cd4⁺ t cells from patients with primary Sjögren's syndrome. Arthritis Rheumatol 2014; 66: 731-9.
- 7. KONSTA OD, THABET Y, LE DANTEC C *et al.*: The contribution of epigenetics in Sjögren's syndrome. *Front Genet* 2014; 5: 71.
- KIVITY S, ARANGO MT, EHRENFELD M et al.: Infection and autoimmunity in Sjögren's syndrome: A clinical study and comprehensive review. J Autoimmun 2014; 51: 17-22.
- 9. BURBELO PD, AMBATIPUDI K, ALEVIZOS I: Genome-wide association studies in Sjögren's syndrome: What do the genes tell us about disease pathogenesis? *Autoimmun Rev* 2014; 7: 756-61.
- TEOS LY, ZHANG Y, COTRIM AP et al.: Ip3r deficit underlies loss of salivary fluid secretion in Sjögren's syndrome. Sci Rep 2015; 13953.

- WYNN TA, RAMALINGAM TR: Mechanisms of fibrosis: Therapeutic translation for fibrotic disease. *Nat Med* 2012; 7: 1028-40.
- GARDET A, ZHENG TS, VINEY JL: Genetic architecture of human fibrotic diseases: Disease risk and disease progression. *Front Pharmacol* 2013; 159.
- HSIEH C, CHANG A, BRANDT D, GUTTIKON-DA R, UTSET TO, CLARK MR: Predicting outcomes of lupus nephritis with tubulointerstitial inflammation and scarring. *Arthritis Care Res* 2011; 6: 865-74.
- BOWLUS CL, GERSHWIN ME: The diagnosis of primary biliary cirrhosis. *Autoimmun Rev* 2014; 4-5: 441-4.
- HUDSON M, FRITZLER MJ: Diagnostic criteria of systemic sclerosis. J Autoimmun 2014; 38-41.
- CONRAD K, ROGGENBUCK D, LAASS MW: Diagnosis and classification of ulcerative colitis. *Autoimmun Rev* 2014; 13: 463-6.
- PETRI M, ORBAI AM, ALARCON GS et al.: Derivation and validation of the systemic lupus international collaborating clinics classification criteria for systemic lupus erythematosus. Arthritis Rheum 2012; 8: 2677-86.
- REICHLIN M, HARLEY JB: Immune response to the RNA protein particles in systemic lupus erythematosus. A distinctive dichotomy. *Am J Med* 1988; 6A: 35-7.
- 19. HIRSCHFIELD GM, INVERNIZZI P: Progress in the genetics of primary biliary cirrhosis. *Semin Liver Dis* 2011; 2: 147-56.
- 20. ASSASSI S, MAYES MD: What does global gene expression profiling tell us about the pathogenesis of systemic sclerosis? *Curr Opin Rheumatol* 2013; 6: 686-91.
- 21. GEREMIA A, BIANCHERI P, ALLAN P, CORAZ-ZA GR, DI SABATINO A: Innate and adaptive immunity in inflammatory bowel disease. *Autoimmun Rev* 2014; 1: 3-10.
- 22. VITALI C, BOMBARDIERI S, JONSSON R et al.: Classification criteria for Sjögren's syndrome: A revised version of the European criteria proposed by the American-European consensus group. Ann Rheum Dis 2002; 6: 554-8.
- 23. TAPINOS NI, POLIHRONIS M, TZIOUFAS AG, SKOPOULI FN: Immunopathology of Sjögren's syndrome. Ann Med Interne 1998; 1: 17-24.
- 24. DISTLER O, PAP T, KOWAL-BIELECKA O et al.: Overexpression of monocyte chemoattractant protein 1 in systemic sclerosis: Role of platelet-derived growth factor and effects on monocyte chemotaxis and collagen synthesis. Arthritis Rheum 2001; 11: 2665-78.
- 25. TAKEDAY: Histopathological studies of the labial salivary glands in patients with Sjögren's syndrome. Part i: Light microscopic study. *Bull Tokyo Med Dent Univ* 1980; 1: 9-25.
- 26. SKOPOULI FN, LI L, BOUMBA D et al.: Association of mast cells with fibrosis and fatty infiltration in the minor salivary glands of patients with Sjögren's syndrome. Clin Exp Rheumatol 1998; 1: 63-5.
- 27. BOOKMAN AA, SHEN H, COOK RJ et al.: Whole stimulated salivary flow: Correlation with the pathology of inflammation and damage in minor salivary gland biopsy specimens from patients with primary Sjögren's syndrome but not patients with sicca. Arthri-

tis Rheum 2011; 7: 2014-20.

- LLAMAS-GUTIERREZ FJ, REYES E, MAR-TINEZ B, HERNANDEZ-MOLINA G: Histopathological environment besides the focus score in Sjögren's syndrome. *Int J Rheum Dis* 2014; 17: 898-903.
- 29. SYRJANEN S: Age-related changes in structure of labial minor salivary glands. *Age Ageing* 1984; 3: 159-65.
- 30. GREENSPAN JS, DANIELS TE, TALAL N, SYLVESTER RA: The histopathology of Sjögren's syndrome in labial salivary gland biopsies. Oral Surg Oral Med Oral Pathol 1974; 2: 217-29.
- 31. DANIELS TE, SILVERMAN JR S, MICHALSKI JP, GREENSPAN JS, SYLVESTER RA, TALAL N: The oral component of Sjögren's syndrome. Oral Surg Oral Med Oral Pathol 1975; 6: 875-85.
- 32. DANIELS TE, WHITCHER JP: Association of patterns of labial salivary gland inflammation with keratoconjunctivitis sicca. Analysis of 618 patients with suspected Sjögren's syndrome. Arthritis Rheum 1994; 6: 869-77.
- 33. TAKEDA Y: Histopathological studies of the labial salivary glands in patients with Sjogren's syndrome. Part ii: Electron microscopic study. *Bull Tokyo Med Dent Univ* 1980; 1: 27-42.
- 34. RASMUSSEN A, ICE JA, LI H et al.: Comparison of the American-European consensus group Sjögren's syndrome classification criteria to newly proposed American College of Rheumatology criteria in a large, carefully characterised sicca cohort. Ann Rheum Dis 2014; 73: 31-8.
- 35. VITALI C, BOMBARDIERI S, MOUTSOPOU-LOS HM *et al.*: Preliminary criteria for the classification of Sjögren's syndrome. Results of a prospective concerted action supported by the european community. *Arthritis Rheum* 1993; 3: 340-7.
- 36. JOACHIMS ML, LEEHAN KM, LAWRENCE C et al.: Single-cell analysis of glandular T cell receptors in Sjögren's syndrome. JCI Insight 2016; 1: pii: e85609.
- 37. R CORE TEAM RFFSC, VIENNA, AUSTRIA R: A language and environment for statistical computing. 2013.
- 38. FOX JAW, SANFORD: An {r} companion to applied regression, second edition. 2011.
- 39. ROBIN X, TURCK N, HAINARD A *et al.*: pROC: An open-source package for R and S+ to analyze and compare ROC curves. *BMC Bioinformatics* 2011; 12: 77.
- 40. LÓPEZ-RATÓN M, RODRÍGUEZ-ÁLVAREZ MX, CADARSO-SUÁREZ C, GUDE-SAMPE-DRO F: Optimalcutpoints: An r package for selecting optimal cutpoints in diagnostic tests. 2014 2014; 8: 36.
- 41. GENUER R, POGGI J-M, TULEAU-MALOT C, GENUER MR: Package 'vsurf'. Pattern Recognition Letters 2015; 14: 2225-36.
- 42. DANIELS TE: Labial salivary gland biopsy in Sjögren's syndrome. Assessment as a diagnostic criterion in 362 suspected cases. *Arthritis Rheum* 1984; 2: 147-56.
- 43. IWANO M, PLIETH D, DANOFF TM, XUE C, OKADA H, NEILSON EG: Evidence that fibroblasts derive from epithelium during tissue fibrosis. *J Clin Invest* 2002; 3: 341-50.
- 44. FRIEDMAN SL, SHEPPARD D, DUFFIELD JS, VIOLETTE S: Therapy for fibrotic diseases:

Salivary gland fibrosis in Sjögren's syndrome / K.M. Leehan et al.

Salivary gland fibrosis in Sjögren's syndrome / K.M. Leehan et al.

Nearing the starting line. *Sci Transl Med* 2013; 5: 167sr1.

- 45. MCARTHUR CP, DANIELS PJ, KRAGEL P, HOWARD PF, JULIAN L: Sjögren's syndrome salivary gland immunopathology: Increased laminin expression precedes lymphocytic infiltration. *J Autoimmun* 1997; 1: 59-65.
- 46. KAPSOGEORGOU EK, DIMITRIOU ID, ABU-HELU RF, MOUTSOPOULOS HM, MANOUS-SAKIS MN: Activation of epithelial and myoepithelial cells in the salivary glands of patients with Sjögren's syndrome: High ex-

pression of intercellular adhesion molecule-1 (icam.1) in biopsy specimens and cultured cells. *Clin Exp Immunol* 2001; 1: 126-33.

- 47. KWON YJ, PEREZ P, AGUILERA S et al.: Involvement of specific laminins and nidogens in the active remodeling of the basal lamina of labial salivary glands from patients with Sjögren's syndrome. Arthritis Rheum 2006; 11: 3465-75.
- LESSARD CJ, LI H, ADRIANTO I et al.: Variants at multiple loci implicated in both innate and adaptive immune responses are associ-

ated with Sjögren's syndrome. *Nat Genet* 2013; 11: 1284-92.

- 49. DIEUDE P, GUEDJ M, WIPFF J *et al.*: Association between the irf5 rs2004640 functional polymorphism and systemic sclerosis: A new perspective for pulmonary fibrosis. *Arthritis Rheum* 2009; 1: 225-33.
- KAPSOGEORGOU EK, CHRISTODOULOU MI, PANAGIOTAKOS DB *et al.*: Minor salivary gland inflammatory lesions in Sjögren's syndrome: Do they evolve? *J Rheumatol* 2013; 9: 1566-71.