

Erratum corrigé

Stressful life events are associated with the diagnosis of systemic autoimmune rheumatic diseases among adults

I. Bauer Ventura, M.E. Goldberg, A. Schiffenbauer, M. Shi, R. Volochayev, S.H. Jackson, A. Jansen, N. Bayat, P. Noroozi Farhadi, C.G. Parks, C.R. Weinberg, A. Picardi, F.W. Miller, L.G. Rider

Published in: Clin Exp Rheumatol 2025; 43(1): 21-27.

The authors have brought to our attention that Table VI was missing from the published article in the printed paper issue.

Table VI. Adjusted conditional logistic regression modelling of life events scores with systemic autoimmune rheumatic disease diagnosis among probands and their unaffected siblings*.

Life event/scores by category	Odds ratio	95% CI	p-value
Number of life events by category:			
Total number of events	1.07	0.94–1.23	0.309
Major events	0.96	0.52–1.78	0.899
Controllable events	1.02	0.59–1.76	0.956
Uncontrollable events	1.23	0.92–1.65	0.165
Desirable events	0.77	0.36–1.64	0.500
Undesirable events	1.12	0.88–1.42	0.369
Highly stressful events	1.12	0.92–1.36	0.270
Weighted event scores by category:			
Weighted total event score	1.04	0.98–1.10	0.211
Weighted major event score	1.05	0.84–1.31	0.678
Weighted controllable event score	1.10	0.78–1.54	0.595
Weighted uncontrollable event score	1.10	0.97–1.24	0.158
Weighted desirable event score	1.02	0.67–1.56	0.911
Weighted undesirable event score	1.05	0.96–1.16	0.304

* Models were adjusted for age, sex, race/ethnicity, educational, cigarette smoking ever, and time from date of systemic rheumatic disease diagnosis in probands (or reference date in unaffected siblings) to date of questionnaire completion.
CI: confidence interval.