

Temporomandibular joint involvement and reduced mouth opening in systemic sclerosis: a not interdependent relationship

Sir,

Temporomandibular joint (TMJ) involvement in rheumatic diseases is often neglected. Temporomandibular symptoms and signs in immune-mediated rheumatic diseases are more frequent in juvenile idiopathic arthritis (30-93%) (1). Systemic sclerosis (SSc) is an autoimmune connective tissue disease, which involves the immune system and microcirculation, leading to fibrosis of the skin, subcutaneous tissue and internal organs. The musculoskeletal system is also affected (2-3). In SSc the mouth opening reduction could depend on joint inflammation, as in other rheumatic diseases, but also on tissue fibrosis which can involve skin, subcutaneous tissues and even articular and periarticular structures (4-6) (Fig. 1a). The main pathologic changes of TMJ may be detected by imaging techniques or directly by arthroscopy (5, 7, 8). In recent years minimally invasive surgery (MIS) as arthroscopic disc repositioning and arthroscopic condyle remodelling has been widely used in several pathologic conditions involving TMJ (9). The evaluation of the mechanisms provoking reduced mouth opening may be very difficult in some SSc patients, as in the case we report. This study was performed in accordance with the Helsinki Declaration of 1975/83. Informed consent was obtained from the patient included in the study.

A seventy-year-old woman with severe SSc who had been suffering for many years was referred to the Rheumatology unit of the University of Florence. Multi-organ involvement was present. The clinical picture was characterised by ulcers of fingers of hands and feet. The patient showed a typical aspect of the scleroderma face with amimia. Gastrointestinal apparatus was mainly involved with dysphagia and pyorrhoia associated to reduced opening of the mouth (7 mm maximum interincisal distance) that did not allow a normal feeding. Severe bilateral TMJ pain was present at rest and during the movement. The extreme mouth opening reduction and the pain did not even allow to perform the oesophagogastrosocopy which was necessary to investigate the characteristics of oesophageal involvement provoking dysphagia and anorexia. In this patient we used ultrasound (US) examination to evaluate whether pain and reduced mouth opening were due to the inflammatory involvement of joint or not and the condyle bone remodelling was evident (Fig. 1b). The patient was consequently referred to an oral and maxillofacial unit. The first therapeutic approach to induce pain relief and to increase mouth opening,




Fig. 1. a) Anatomical TMJ features with position of external capsular ligament; b) TMJ capsular width US image as shown by callipers without joint effusion; c) region of interest for V-MIS; d), e), f) sequence of steps in V-MIS; g) mouth opening before surgical treatment; h) mouth opening after surgical treatment.

was the local injection with corticosteroids and hyaluronic acid without any effects. Considering the absence of inflammatory signs involving both TMJs, it was hypothesised that reduced mouth opening could be due to the stiffness of joint capsule and/or periarticular tissues. A videoscope-assisted minimally invasive surgery (V-MIS) was performed under local anaesthesia in an outpatient surgical centre. A bilateral capsular arthrotomy was performed with section of external lateral capsular ligament (Fig. 1c, d, e, f). Immediately after the procedure a wider mouth opening was ob-

tained (13 mm) and pain disappeared. In follow up, 10 days after V-MIS, the maximum interincisal distance was increased from 13 to 18 mm, and the absence of pain was persistent (Fig. 1g, h). After one month the patient still reported no TMJ pain at rest and during mouth opening and had regular feeding. The increase of mouth opening allowed oesophagogastrosocopy.

In this difficult management a multidisciplinary evaluation is mandatory for a tailored therapeutic approach. The discussion of the case in a multidisciplinary team allowed us to obtain a shared hypothesis. It is

Letters to the Editors

to note that the hypothesis *ad excludendum* of periarticular fibrosis was considered, also evaluating the good therapeutic effect obtained after surgical section of capsular ligaments.

M.A. BAGNI, *PhD*
S. GUIDUCCI, *MD*
G. LEPRI, *MD*
S. TORRACCHI, *MD*
P. BERNARDINI, *MD*
D. MELCHIORRE, *PhD*

Department of Experimental and Clinical Medicine, University of Florence, Italy.

Please address correspondence to:

Daniela Melchiorre

Department of Experimental and Clinical Medicine,

Division of Rheumatology,

University of Florence,

Largo Brambilla 3, 50134 Florence, Italy.

E-mail: daniela.melchiorre@unifi.it

ORCID: 0000-0002-8510-0304

Competing interests: none declared.

© Copyright CLINICAL AND
EXPERIMENTAL RHEUMATOLOGY 2026.

References

1. UPADHYAY J, LEMME J, CAY M *et al.*: A multidisciplinary assessment of pain in juvenile idiopathic arthritis. *Semin Arthritis Rheum* 2021; 51(4): 700-11. <https://doi.org/10.1016/j.semarthrit.2021.05.011>
2. LEPRI G, DI BATTISTA M, CODULLO V *et al.*: Systemic sclerosis: one year in review 2024. *Clin Exp Rheumatol* 2024; 42(8): 1517-28. <https://doi.org/10.55563/clinexprheumatol/is29he>
3. CONNOLLY CM, PAIK JJ: Myopathy in systemic sclerosis. *Curr Opin Rheumatol* 2023; 35(6): 341-8. <https://doi.org/10.1097/BOR.0000000000000966>
4. CRINCOLI V, FATONE L, FANELLI M *et al.*: Orofacial manifestations and temporomandibular disorders of systemic scleroderma: an observational study. *Int J Mol Sci* 2016; 17 (7): 1189. <https://doi.org/10.3390/ijms17071189>
5. MATARESE G, ISOLA G, ALIBRANDI A *et al.*: Occlusal and MRI characterization in systemic sclerosis patients: a prospective study from Southern Italian cohort. *Joint Bone Spine* 2016; 83 (1): 57-62. <https://doi.org/10.1016/j.jbspin.2015.04.014>
6. MADDALI BONGI S, PASSALACQUA M, LANDI G *et al.*: Rehabilitation of the face and temporomandibular joint in systemic sclerosis. *Ther Adv Musculoskel Dis* 2021; 13: 1-10. <https://doi.org/10.1177/1759720x211020171>
7. HYSAE, LERCARAA, CERE A *et al.*: Temporomandibular disorders in immune-mediated rheumatic diseases of the adult: a systematic review. *Semin Arthritis Rheum* 2023; 61: 152215. <https://doi.org/10.1016/j.semarthrit.2023.152215>
8. GUIDUCCI S, BAGNI MA, TORRACCHI S *et al.*: An ultrasound working table as diagnostic tool of temporomandibular joint inflammation. *J Ultrasound* 2025; 28 (1): 129-36. <https://doi.org/10.1007/s40477-025-00990-4>
9. ÂNGELO DF, LOPES CS, SANZ D *et al.*: Temporomandibular joint minimally invasive procedures in pediatric population: a prospective study. *J Clin Med* 2024; 13: 672. <https://doi.org/10.3390/jcm13030672>