

# Clinical features, treatment and long-term outcomes of Schnitzler syndrome: a report from China

Sir,  
Schnitzler syndrome (SchS) is a rare acquired autoinflammatory disorder characterised by chronic urticarial rash, monoclonal gammopathy and systemic inflammation, diagnosed by the Strasbourg criteria (1). Most reported cohorts are European; data from China are scarce, and the prolonged unavailability of interleukin-1 (IL-1) inhibitors has restricted access to first-line therapy. We report eight consecutive patients fulfilling these criteria at Peking Union Medical College Hospital (2018-2025), the largest Chinese series to date. The study has been approved by the Institutional Review Board of Peking Union Medical College Hospital. All methods followed the principles outlined in the Declaration of Helsinki for human. Written and informed consent was obtained from all participants. Demographic, clinical, laboratory, treatment and outcome data were retrieved retrospectively from the medical records. Genetic testing was performed by whole-exome sequencing (WES) of genomic DNA from peripheral blood (n=4) or bone marrow (n=2). Treatment response was assessed at each follow-up visit, usually at six-month intervals, and categorized as complete, partial, or no response based on clinical remission and normalization of inflammatory markers by more than 80%, 20%-80%, or less than 20%, respectively. All eight patients were Han Chinese and seven were male. Median age at onset was 52 years and median diagnostic delay 5.5 years. All had chronic urticaria; fever oc-

curred in seven, arthralgia or bone pain in seven, lymphadenopathy in five and hearing loss in one. In four patients, fever developed a median of 2 years after rash onset. All had leukocytosis and raised inflammatory markers (median CRP 119 mg/L, ESR 95 mm/h). Monoclonal gammopathy was IgM- $\kappa$  in seven and IgM- $\lambda$  in one. Neutrophilic urticarial dermatosis was demonstrated in five patients and bone-imaging abnormalities in six. One patient carried an *NLRP3* c.386A>G variant of uncertain significance; none of the six patients tested carried *MYD88* L265P, reported in approximately 30% of patients with SchS(2). Only one patient received an IL-1 inhibitor (canakinumab, obtained overseas) and achieved complete remission; four received tocilizumab with partial responses. Conventional agents and off-label Janus kinase (JAK) or Bruton tyrosine kinase (BTK) inhibitors gave partial benefit only (Table I). Over a mean follow-up of 14 months (range 3-24), one patient developed Waldenström macroglobulinaemia (WM). Within the limits of a small, single-centre series, our patients showed a male predominance (7:1) and exclusively IgM gammopathy, more pronounced than in Western cohorts (male-to-female ratio 1.2-1.8:1; IgG in 7-15%) (3, 4). This may reflect the small sample and referral bias rather than true ethnic variation and requires confirmation in larger cohorts. The absence of *MYD88* L265P warrants particular caution: only six patients were tested, not including the one who developed WM, and the mutation in SchS is largely B-cell-restricted and, without B-cell enrichment, is readily missed by WES, whereas the reported prevalence of around 30% derives from targeted allele-specific PCR with a sensitivity of about 1% (2). Our negative result therefore probably

reflects assay sensitivity rather than true absence, and targeted allele-specific or droplet-digital PCR on purified B cells would be preferable in future studies. The most striking finding is the treatment gap: only one of eight patients received IL-1 blockade, far below utilisation in Western practice (5), reflecting drug availability rather than clinical indication. Where IL-1 inhibitors were inaccessible, IL-6 blockade (6) and JAK or BTK inhibitors (7, 8) achieved partial control only, consistent with previous Chinese reports (9). The short follow-up also limits assessment of long-term complications: SchS carries an estimated 15% 10-year risk of WM, a 15-20% risk of haematological malignancy and a risk of AA amyloidosis (10). These outcomes cannot yet be captured here and mandate prolonged surveillance. In conclusion, this Chinese series confirms the classic SchS phenotype while exposing a critical shortfall in access to first-line IL-1 inhibitors. Expanded access, evaluation of alternatives such as IL-6, JAK and BTK inhibitors, comprehensive genetic testing and long-term monitoring are urgently needed.

S. WANG<sup>1</sup>, MD  
C. LI<sup>1</sup>, MD  
D. WU<sup>2</sup>, MD  
M. SHEN<sup>1,2</sup>, MD

<sup>1</sup>Department of Rare Diseases, Peking Union Medical College Hospital (PUMCH), Chinese Academy of Medical Sciences & Peking Union Medical College; State Key Laboratory of Complex Severe and Rare Diseases, PUMCH, Beijing, China.

<sup>2</sup>Department of Rheumatology and Clinical Immunology, PUMCH; National Clinical Research Center for Dermatologic and Immunologic Diseases (NCRC-DID), Ministry of Science & Technology; Key Laboratory of Rheumatology and Clinical Immunology, Ministry of Education, Beijing, China.

**Table I.** Genetic, therapeutic and outcome data of eight patients with Schnitzler syndrome.

| Pt | Sex/age at onset (y) | Delay (y) | M-protein      | Genetic testing                          | Key clinical features                           | Post-diagnosis targeted therapy (dose)                               | Follow-up (mo.) | Response |
|----|----------------------|-----------|----------------|------------------------------------------|-------------------------------------------------|----------------------------------------------------------------------|-----------------|----------|
| 1  | M/55                 | 11        | IgM- $\kappa$  | WES (blood): negative                    | Rash, fever, LN, hearing loss                   | Tocilizumab 400 mg q4w + prednisone + MTX                            | 18              | Partial  |
| 2  | M/53                 | 5         | IgM- $\lambda$ | WES (blood): negative                    | Rash, fever, arthralgia, LN                     | Tocilizumab 560 mg q4w + prednisone + HCQ → canakinumab 150 mg q8w   | 24              | Complete |
| 3  | F/35                 | 5         | IgM- $\kappa$  | WES (blood): negative                    | Rash, fever, arthralgia                         | Prednisone + Tripterygium wilfordii glycosides                       | 17              | Partial  |
| 4  | M/52                 | 7         | IgM- $\kappa$  | WES (bone marrow): negative              | Rash, fever, arthralgia, LN                     | Tocilizumab 400 mg q4w + prednisone + colchicine + tofacitinib + HCQ | 18              | Partial  |
| 5  | M/44                 | 1         | IgM- $\kappa$  | Not performed                            | Rash, fever, arthralgia, LN                     | Zanubrutinib + prednisone + thalidomide                              | 6               | Partial  |
| 6  | M/60                 | 6         | IgM- $\kappa$  | WES (bone marrow): negative              | Rash, fever, arthralgia, LN                     | Tofacitinib + prednisone + colchicine + thalidomide                  | 3               | Partial  |
| 7  | M/54                 | 8         | IgM- $\kappa$  | Not performed                            | Rash, arthralgia; developed WM during follow-up | Prednisone + Tripterygium wilfordii glycosides                       | 10              | Partial  |
| 8  | M/62                 | 0.1       | IgM- $\kappa$  | WES (blood): <i>NLRP3</i> c.386A>G (VUS) | Rash, fever, arthralgia                         | Tocilizumab 400 mg, irregular intervals + prednisone                 | 16              | Partial  |

Pt: patient; y: years; LN: lymphadenopathy; WES: whole-exome sequencing; VUS: variant of uncertain significance; MTX: methotrexate; HCQ: hydroxychloroquine; WM: Waldenström macroglobulinaemia; q4w/q8w: every 4/8 weeks.

# Letters to the Editors

Please address correspondence to:

Min Shen

Department of Rare Diseases,

Peking Union Medical College Hospital

(PUMCH),

Chinese Academy of Medical Sciences &

Peking Union Medical College,

State Key Laboratory of Complex Severe and

Rare Diseases, PUMCH,

Beijing 100730, China.

E-mail: shenmpumch@163.com

**Funding.** CAMS Innovation Fund for Medical

Sciences (CIFMS) (2024-I2M-C&T-B-014);

National High Level Hospital Clinical

Research Funding (2025-PUMCH-D-006);

Open Research Fund of the State Key

Laboratory of Complex, Severe, and Rare

Diseases (2025-O-ZD-005).

**Competing interests:** none declared.

## References

1. SIMON A, ASLI B, BRAUN-FALCO M *et al.*: Schnitzler's syndrome: Diagnosis, treatment, and follow-up. *Allergy* 2013; 68(5): 562-8. <https://doi.org/10.1111/all.12129>
2. PATHAK S, ROWCZENIO DM, OWEN RG *et al.*: Exploratory study of myd88 1265p, rare nlrp3 variants, and clonal hematopoiesis prevalence in patients with schnitzler syndrome. *Arthritis Rheumatol* 2019; 71(12): 2121-5. <https://doi.org/10.1002/art.41030>
3. ROWCZENIO DM, PATHAK S, AROSTEGUI JI *et al.*: Molecular genetic investigation, clinical features, and response to treatment in 21 patients with schnitzler syndrome. *Blood* 2017; 131(9): 974-81. <https://doi.org/10.1182/blood-2017-10-810366>
4. DE KONING HD: Schnitzler's syndrome: Lessons from 281 cases. *Clin Transl Allergy* 2014; 4: 41. <https://doi.org/10.1186/2045-7022-4-41>
5. CALABRESE L, CARTOCCI A, VITALE A *et al.*: IL-1 targeting agents in schnitzler syndrome: A multicentre, real-world study from the international aida network schnitzler registry. *Clin Exp Rheumatol* 2025; 43(10): 1753-62. <https://doi.org/10.55563/clinexprheumatol/i9hsfr>
6. BONNEKOH H, FRISCHBUTTER S, ROLL S, MAURER M, KRAUSE K: Tocilizumab treatment in patients with schnitzler syndrome: An open-label study. *J Allergy Clin Immunol Pract* 2021; 9(6): 2486-9. <https://doi.org/10.1016/j.jaip.2021.01.024>
7. LI M, CHEN Y-W, SHEN A *et al.*: Case report: Successful treatment with tofacitinib and colchicine in a patient with schnitzler syndrome. *Int J Rheum Dis* 2022; 26(1): 160-3. <https://doi.org/10.1111/1756-185X.14457>
8. HUANG Y, WANG Y, YU F *et al.*: Case report: Therapeutic use of ibrutinib in a patient with schnitzler syndrome. *Front Immunol* 2022; 13: 894464. <https://doi.org/10.3389/fimmu.2022.894464>
9. YAN R, CAO W, LIU X, LI F, SHEN M: A chinese case series of schnitzler syndrome and complete remission in one tocilizumab-treated patient. *Clin Rheumatol* 2020; 39(12): 3847-52. <https://doi.org/10.1007/s10067-020-05204-2>
10. LIPSKER D: The schnitzler syndrome. *Orphanet J Rare Dis* 2010; 5: 38. <https://doi.org/10.1186/1750-1172-5-38>